		Type: Drug Guideline	Document reference: 0569	Manual Classification: Waikato DHB Drug Guidelines
Title: Amoxicillin for Neonates			Effective date: 24 August 2020	
Facilitator <small>sign/date</small>	Authorised <small>sign/date</small>	Authorised <small>sign/date</small>	Version: 3	Page: 1 of 2
<i>Kerrie Knox Pharmacist</i>	<i>Jutta van den Boom Clinical Director NICU</i>	<i>John Barnard Chair Medicines & Therapeutics</i>	Document expiry date: 24 August 2023	

© Waikato DHB, September 2020

BRIEF ADMINISTRATION GUIDE

For detailed information refer to [The Australasian Neonatal Medicines Formulary amoxicillin guideline](#)



Critical Note: there are minor variations between the ANMF and Waikato DHB best practice within this drug guideline – see **yellow shaded text**

- Indications:**
- Bacterial infections sensitive (or likely to be) to amoxicillin
 - Early sepsis (≤ 2 days) empiric therapy (in combination with gentamicin)

Route: Intravenous, intramuscular, oral

- Injection supplied as amoxicillin sodium, equivalent to amoxicillin 250 mg vial
- Oral (**request from Pharmacy**) supplied as amoxicillin 250 mg/5mL or 125 mg/5mL suspension
- pH of amoxicillin 8 to 10

Dose:

Route	Dose (mg/kg/dose)	Postnatal age (days)	Dosing Interval (hours)
IV	50*	0 to 7	12
		>7	8

*Note: dose at 100 mg/kg/dose IV for severe infections e.g. meningitis

Oral dosing for non-systemic infections: 30 mg/kg (maximum 125 mg) three times daily

Preparation and administration

Intravenous

- Reconstitute **250 mg vial** with **4.8 mL** of water for injection to make a concentration of **50 mg/mL** (Note: 250 mg of amoxicillin displaces 0.2 mL of diluent)
- Shake vial vigorously, as soon as diluent is added, to dissolve powder and check for absence of particulate matter before proceeding
- Draw up the required volume for the dose and if necessary dilute further with sodium chloride 0.9%
- Infuse dose over **30 minutes** using Guardrails profiled syringe driver
- Flush before and after the dose with sodium chloride 0.9% (preferably) or **glucose 5 or 10 %**.
Note: although amoxicillin is reported to be unstable with glucose (the medication degrades faster) it is deemed acceptable to flush with or administer concurrently when necessary. The flush should be administered at the same rate as the amoxicillin.

Intramuscular


- Dilute **250 mg** vial with **1.3 mL** of water for injection. This produces a final concentration of 167 mg/mL (250mg/1.5mL)
- Note: IM dose volume should be kept to ≤ 1 mL if possible to decrease pain.
- Inject dose into a large muscle e.g. buttock, thigh

Oral

- Shake suspension well then draw up appropriate volume in an oral syringe. The dose may be mixed with milk or water if required. After mixing administer immediately.

Monitoring

- Consider if specimen for culture and sensitivity testing is required before first dose
- Assess for signs of anaphylaxis and adverse reactions
- Observe infusion site for thrombophlebitis
- Monitor temperature and other parameters appropriate to the condition

	Document reference: 0569	Effective date: 24 Aug 2020	Expiry date: 24 Aug 2023	Page: 2 of 2
	Title: Amoxicillin for neonates	Type: Drug Guideline	Version: 3	Authorising initials:

- Observe for change in bowel frequency
- Monitor renal, hepatic and hematologic function periodically

Storage and Stability

- Reconstituted solution in the vial and solutions diluted further with sodium chloride 0.9% are stable for **up to 6 hours**.
- The reconstituted oral suspension is stable for 14 days at room temperature or refrigerated (preferred).

Competency for Administration

This procedure is carried out by, or under, the direct supervision of a registered nurse/registered midwife who holds current Waikato DHB Generic Medicine Management and IV certification plus Guardrails competency (if administering IV) as well as Neonatal specific competency NCV/NAC (if administering via CVAD).

Guardrails Information

Amoxicillin is Guardrail profiled on the CC pump for NICU. Following are the Guardrail limits:

Guardrails Drug Name Pump	Amoxicillin*		
	0.4-1kg	1-2kg	2-3 & 3-5kg
Concentration (mg/ml)			
Minimum	12.5	31	50
Maximum	100	100	100
Dose rate (mg/kg/hour)			
Default	100	100	100
Soft minimum	50	50	50
Soft maximum	200	200	200
Hard max	202	202	202

References

- Royal Prince Alfred Hospital NeoMed Consensus Group. Amoxicillin Drug Information Sheet. October 2019. Available from: <https://www.slhd.nsw.gov.au/rpa/neonatal/NeoMedPaperCopy.html>
- Douglas Pharmaceuticals Ltd. Ibiamox® Data Sheet. 11 April 2018. Available from <https://www.medsafe.govt.nz/profs/datasheet/i/ibiamoxinj.pdf>
- Truven Health Analytics Inc. Pediatrics and Neofax®. 2019. Amoxicillin monograph. Accessed 7.12.2019. Available from: <http://www.micromedexsolutions.com>.
- New Zealand Formulary for Children (NZFC). Amoxicillin. Accessed 7.12.2019. Available from https://nzfchildren.org.nz/nzf_3025
- The Royal Children's Hospital Melbourne. Paediatric Injectable Guidelines. Accessed 7.12.2019. Available from <https://pig.rch.org.au>.
- New Zealand Hospital Pharmacists Assoc. Inc. Notes on Injectable Drugs 7th ed, 2015.
- Auckland DHB Newborn Services. Amoxycillin Drug Protocol. August 2016. Available from <http://www.adhb.govt.nz/newborn/DrugProtocols/AmoxycillinPharmacology.htm>
- Canterbury DHB Neonatal Services. Amoxycillin (Parenteral) Drug Information Sheet. March 2016. Available from <https://cdhb.health.nz/wp-content/uploads/8ffdfccc-amoxycillin20parenteral.pdf>
- Canterbury DHB Neonatal Services. Amoxycillin (Oral) Drug Information Sheet. March 2016. Available from <https://cdhb.health.nz/wp-content/uploads/a0a5f31c-amoxycillin20oral.pdf>

Note: Printed copies are only valid on the day of printing – they are not controlled and may not be the current version in use. Please refer to the online version.

Disclaimer: This document has been developed by Waikato District Health Board specifically for its own use. Use of this document and any reliance on the information contained therein by any third party is at their own risk and Waikato District Health Board assumes no responsibility whatsoever.